

Global scale evidence for the refractory nature of riverine black carbon

Alysha I. Coppola¹, Daniel B. Wiedemeier¹, Valier Galy⁵, Negar Haghipour², Ulrich M. Hanke¹, Gabriela S. Nascimento², Muhammed Usman², Thomas M. Blattmann², Moritz Reisser¹, Chantal V. Freymond², Meixun Zhao⁴, Britta Voss⁵, Lukas Wacker³, Enno Schefuß⁶, Bernhard Peucker-Ehrenbrink⁵, Samuel Abiven¹, Michael W.I. Schmidt¹, Timothy I. Eglinton^{2,5}

*Corresponding author, email: Alysha.coppola@geo.uzh.ch o

¹ Department of Geography, University of Zurich, Winterthurerstrasse 190, 8057 Zürich Switzerland

²Geological Institute, Department of Earth Sciences, ETH Zürich, Sonneggstrasse 5, 8092 Zürich Switzerland

³Laboratory of Ion Beam Physics, ETH Zürich, Otto-Stern-Weg 5, 8093 Zürich, Switzerland

⁴Key Laboratory of Marine Chemistry Theory and Technology of the Ministry of Education, Ocean University of China, 238 Songling Road Qingdao 266100/Laboratory for Marine Ecology and Environmental Science, Qingdao National Laboratory for Marine Science and Technology, Qingdao 266061, China

⁵Woods Hole Oceanographic Institution, Department of Marine Chemistry and Geochemistry, 360 Woods Hole Road, Woods Hole, Massachusetts 02543 USA

⁶MARUM-Center for Marine Environmental Sciences, University of Bremen, Bremen, Germany

24

25 **ABSTRACT**

26 Wildfires and fossil fuel combustion release large amounts of greenhouse gases into
27 the atmosphere, but also produce Black Carbon (BC, or pyrogenic carbon) from
28 incomplete combustion. Processes controlling BC production and its fate are an
29 integral component of the carbon cycle. Constraining BC export from land to the
30 ocean is critical given on-going changes in land use and climate that affect fire
31 occurrence and BC dynamics. Here, we compile an inventory of concentration and
32 radiocarbon contents ($\Delta^{14}\text{C}$) of particulate BC (PBC) for a globally distributed suite of
33 rivers, and show that PBC fluxes co-vary with river sediment particulate organic carbon
34 (POC), indicating that PBC export is primarily controlled by erosion. River PBC is not
35 exclusively from modern sources but includes PBC that has aged (up to $17,000 \pm 780$
36 ^{14}C yrs) from intermediate terrestrial carbon pools in several high latitude rivers. The
37 global, flux-weighted ^{14}C age of PBC delivered to the ocean ($3,700 \pm 400$ ^{14}C yrs, $\Delta^{14}\text{C}$
38 $= -372 \pm 28\%$) implies protracted storage in terrestrial reservoirs before export. River
39 PBC accounts for $15.8 \pm 0.9\%$ of POC, amounting to a global river PBC flux of 0.017 -
40 0.037 Pg yr^{-1} to the oceans. This corresponds to 4-32% of the of global annual BC
41 production, implying an export efficiency that is one to two orders of magnitude greater
42 than for POC. When buried in marine sediments, PBC is sequestered, forming an
43 important long-term sink for atmospheric CO_2 .

44

45 Forest fires and fossil fuel combustion release large amounts of carbon as greenhouse
46 gases and aerosols into the atmosphere, contributing to the on-going changes in
47 Earth's climate that are occurring at an unprecedented rate ¹. Up to 27% of this fire-
48 derived carbon is transformed into Black Carbon (BC, or pyrogenic carbon, charcoal

residues ²⁾ – a byproduct of incomplete combustion - rather than emitted as greenhouse gases ³⁾. The majority of BC is from vegetation fires (Figure 1). Once incorporated into surface reservoirs (e.g. soils, lake sediments), BC participates in many biogeochemical processes, and influences carbon cycling on local to global scales ^{3,4)}. Biomass burning transfers carbon from fast-cycling (atmosphere-biosphere) pools to more slowly cycling soil and sedimentary reservoirs ⁵⁾, creating a long-term carbon sink ^{6,7)}. Due to its aromatic structure, a substantial fraction of BC decomposes slowly ^{8,9)}, and can persist in soils for hundreds to thousands of years ^{5,10)}.

Greater understanding of the role of this slow-cycling component of the carbon cycle and its significance as a sink of atmospheric CO₂ requires improved constraints on the origin, dynamics and fate of BC. River systems connect terrestrial and marine carbon cycles, exporting approximately 2.7 Pg C yr⁻¹ to the oceans ¹¹⁾, where it is ultimately either mineralized to CO₂ and CH₄ ¹²⁻¹⁴⁾ or sequestered in sediments ¹⁵⁾. Rivers deliver BC from land to the ocean both as particulate BC (PBC) in particulate organic carbon (<63 µm) and dissolved BC (DBC) in dissolved organic carbon (<1 µm)^{4,16)}. Dissolved BC, which comprises a substantial fraction (10%) of dissolved organic carbon (OC) globally, is continuously exported from soils for decades after wildfire burning ^{17,18)} (26.5 Tg yr⁻¹), and can cycle in the deep ocean on millennial timescales (~ 20,000 ¹⁴C yrs) ^{19,20)}. The global amount and age of PBC transported by rivers, has remained largely unknown until now. PBC river fluxes, age and transport is essential for constraining land-ocean transfer as well as assessing its significance as a CO₂ sink by sequestration in continental margin sediments ²¹⁾. Current global PBC flux estimates vary by a factor of 20 (0.005-0.108 Pg yr⁻¹) ^{22,23)}, and the magnitude and timescales of transport, transformation and degradation processes are not well understood ^{16,24)}. As

river basins are facing anthropogenic pressures, both directly via changes to the land surface (e.g. increasing fire intensity and frequency, accelerated deforestation, conversion to agriculture) and fluvial networks (e.g. dams, channelization, irrigation), and indirectly via climate change (e.g. increasing temperatures and an invigorated hydrological cycle)^{11,25,26}, we need to constrain PBC river export to assessing past and future perturbations of this slowly cycling pool in the carbon cycle.

Here, we use the abundance and radiocarbon (¹⁴C) content to constrain the flux and age of river PBC of 18 globally distributed rivers. We quantify river PBC in suspended or deposited sediments collected at the terminus of 11 of the largest rivers worldwide, (such as the Amazon, Congo, Brahmaputra, and major Arctic rivers) and 7 small mountainous rivers²⁷ (S.Table 1). These samples represent 15-34% of the global organic carbon exported by rivers (high and low estimates of export from²⁸). We used river suspended sediments (collected by filtration) or freshly deposited river sediments (<63 μm) (Supplementary Materials, S.Figure 1). We measured BC in Particulate Organic Carbon (POC, as <63um size fraction), providing the age of PBC at or near the river terminus (S.Table 1, S.Figure 1). To characterize PBC derived predominantly from residues of biomass burning, we use chemical oxidation to liberate corresponding benzene polycarboxylic acids (BPCAs)²⁹. The analytical window captured by this method implies that the mass weighted PBC fluxes are conservative under-estimates for PBC fluxes, since it does not include by-products of low-temperature fires (e.g. levoglucosan^{21,30}). We assume BC molecular markers have the same ¹⁴C age within BC. Subsequently, we purified BPCA marker compounds²⁹ and converted them to CO₂ followed by ¹⁴C measurement by gas ion source Accelerator Mass Spectrometry³¹. PBC fluxes from each river were estimated by multiplying the relative PBC

concentration (from BPCA concentrations) with the reported biospheric OC yield of each river given by Galy et al.²⁸, thereby normalizing PBC to biospheric OC export. The biospheric OC yield was calculated using previous reported data and linear correlations between biospheric OC yield and sediment yield in Galy et al.,²⁸ (Supplementary Information).

BC as a major component of riverine particulate organic carbon

We find PBC comprises a significant component of river biospheric POC ($15.8 \pm 0.9\%$ PBC river flux-weighted mean, $n=18$, Supplementary Materials). This proportion of river PBC is similar to global averages of the riverine dissolved BC ($10.6 \pm 0.7\%$)¹⁸ and of soil organic carbon (13.7%)³². The proportion of PBC as a component of river POC ranges from $2.7 \pm 0.4\%$ PBC (Pettaquamscutt) to $32.9 \pm 2.9\%$ PBC (Eel) (Supplementary Materials, S.Table 2, S.Figure 2). Corresponding PBC fluxes from rivers range from 8 ± 2 (Fraser) to 1162 ± 218 Gg yr⁻¹ (Amazon). There is no correlation between BC concentration and river basin drainage size (S.Figure 3), however PBC yield is positively correlated with suspended sediment yield (sediment discharge normalized to the drainage area). This correlation follows a power-law relationship ($r^2=0.61$) (Figure 3a) and indicates that the rate of PBC export is controlled primarily through soil erosion, mobilization and transport processes, much like export of POC^{28,33}. PBC concentrations vary by a half order of magnitude, while suspended sediment yield varies by four orders of magnitude, illustrating that PBC export is primarily controlled by erosion (not concentration). Given that rivers sequester the majority of terrestrial exported POC by burial in ocean sediments³⁴, these observations establish a direct link between soil erosion and PBC sequestration on continental margins.

124

125 **Time-lags between production and riverine export of PBC**

126 We find a large range in PBC $\Delta^{14}\text{C}$ values, indicating fast ($+74\pm 62\text{‰}$, modern, Congo)
127 to slow ($-880\pm 12\text{‰}$, $17,000\pm 780\text{ }^{14}\text{C}$ yrs, Colville) PBC cycling within individual
128 watersheds (Figure 3, S. Table 1). The global flux-weighted age average of $3,700\pm 400$
129 ^{14}C yrs ($-372\pm 28\text{‰}$) is significantly higher (older) than the few measurements on river
130 dissolved BC ($450\pm 280\text{ }^{14}\text{C}$ yrs, $475\pm 150\text{ }^{14}\text{C}$ yrs and $1140\text{ }^{14}\text{C}$ yrs, from ^{16,19,20}).
131 Globally, PBC is older than total POC, with two exceptions in the Godavari and Congo
132 Rivers where PBC is modern (Figure 3). Assuming all BC produced from burning of
133 modern biomass has a mean post-bomb $\Delta^{14}\text{C}$ value of $+100\text{‰}$ and BC derived from
134 fossil fuel combustion has a $\Delta^{14}\text{C}$ value of -1000‰ (i.e., is radiocarbon-depleted), we
135 estimate that $44\pm 28\%$ of river PBC is from fossil fuel contributions (Supplementary
136 Materials). However, assuming only two end members is overly simplistic given the
137 range of potential PBC sources and transport pathways ^{16,35}. A regional study in the
138 Pettaquamscutt River basin (USA) estimated a maximum fossil fuel BC contribution of
139 only 19%, and indicated that there is a time lag between production and river export
140 of PBC³⁵. This implies that BC can “pre-age” during temporary storage in intermediate
141 terrestrial reservoirs (e.g. soils) ³⁵.

142

143 River PBC thus reflects at least three pools: i) biomass-derived char from recent
144 vegetation fires, ii) pre-aged BC (held within the catchment in soils, wetlands and
145 floodplains before river transport and ocean deposition ³⁵), and iii) fossil fuel-derived
146 BC. Fossil fuel-derived BC mass contributions are considered minor for the following
147 reasons. First, the annual production of biomass-derived BC ($114\text{--}383\text{ Tg yr}^{-1}$) is one
148 to two orders of magnitude higher than BC produced by fossil fuel combustion ($2\text{--}29$

Tg yr⁻¹) (Figure 1). Second, soil formation rates span centuries to millennia, and the vast majority of BC eroded from soils must pre-date the beginning of the industrial revolution. Third, there is mounting evidence that soil OC inputs dominate the POC load in most river systems ^{28,36}, as indicated for PBC by our own data. PBC contributions to river PBC pools are therefore likely dominated by inputs from biomass burning.

Given the importance of soil OC as a component of riverine POC export, rivers constitute a source of pre-aged PBC. PBC can be temporarily stored in soils and alluvial deposits for thousands of years ^{37,38}, much like other molecular markers of terrestrial vegetation (e.g., higher plant-derived long-chain *n*-alkanes and *n*-alkanoic acids) ³⁹⁻⁴¹. River PBC ¹⁴C ages are generally older than those of other terrestrial vegetation molecular markers ³⁹⁻⁴¹ implying slower PBC turnover rates. Moreover, the correlation between $\Delta^{14}\text{C}$ values of PBC and POC (Figure 3b, $r^2=0.48$, $p=0.005$) suggests that pre-aged soil OC, including PBC, is an important component of overall OC export. This linear relationship is close to unity (1.05 ± 0.02), indicating that common mechanisms are responsible for aging of both soil OC and BC pools, as well as implying that the ratio of BC and non-BC reactivity is roughly constant, regardless of environmental conditions. PBC experiences pre-aging (relative to POC) in river basins by a relatively constant amount globally, including locations where environmental conditions both favor preservation or mineralization.

Global implications and future outlook

Although sequestration in marine sediments is considered to be the ultimate fate of BC ²¹ (Figure 1), marine sediment BC burial fluxes only account for 3-10% of global

BC production fluxes⁴². In soils, BC constitutes on average 13.7% (ranging to 50%) of total organic carbon³². This inconsistency between BC in soils and marine sediments raises the question as to where the majority of BC produced annually on land goes. Inland waters are both significant holding pools of organic carbon (with storage of up to 50% of carbon along the river-to-ocean continuum), and processors of this carbon^{12,14,43}. In this way, PBC may be sequestered for years up to millennia in intermediate reservoirs prior to export and burial in marine sediments^{35,44}. For example, PBC has been found to be retained in alluvial deposits for thousands of years³⁷, implying that BC can undergo pre-aging en-route to its ultimate burial site. The large fluxes and diverse ages of PBC in our study supports the conclusion that dissolved and particulate BC pools in rivers are partially decoupled, with dissolved BC driven by hydrology^{16,19,45} whereas PBC is subject to erosional and depositional dynamics (Figure 1). This preliminary global assessment of river PBC flux represents an underestimate since it is based on BPCA markers tracing the most refractory components of BC (i.e., it does not include more labile BC from low temperature fires³⁰). Our study highlights the need for further source-to-sink studies to determine controls on the fluxes and degrees of pre-aging of PBC in river catchments prior to export.

Assuming a soil BC stock of 200 Pg³², and a BC production rate of 0.123-0.56 Pg C per yr, the BC mean turnover time in soils at steady state ranges from 1600 to 3500 yrs (Figure 3)⁴. This is much longer than estimates of bulk soil OC turnover times (mean residence time, 25-110 yrs)⁴⁶, highlighting the refractory nature of BC, and consistent with our estimated watershed-wide erosion rates. Using our weighted mean PBC as a fraction of biospheric POC (15.8±0.9%) and estimated global biospheric

POC flux (0.157^{+74}_{-50} Pg yr⁻¹ ²⁸), we estimate an annual global flux of PBC to the ocean of 0.017-0.037 Pg BC. This riverine PBC flux is approximately equal to the global dissolved BC flux (0.027 Pg yr⁻¹)¹⁸ indicating that, relative to atmospheric fluxes (0.002 - 0.005 Pg yr⁻¹)⁴, river transport serves as the dominant process for mobilization of BC from land to ocean (Figure 1). Thus, rivers transport 4-32% of the 0.114-0.383 Pg BC that is produced annually to the oceans. This is one to two orders of magnitude larger than the fraction of biospheric OC that is exported, indicating that a much larger fraction of PBC (relative to biospheric OC or dissolved BC) is laterally exported instead of respired in soils. Furthermore, the average age of riverine PBC ($3,700 \pm 400$ ¹⁴C yrs) is closer to the calculated mean BC turnover time (800-1000 yrs) ⁴⁷ in soils than that of the biospheric OC (~50 yrs)⁴⁸. Together these observations provide global evidence that PBC is more refractory than POC. Like POC ⁴⁹, river PBC is likely to be transferred and buried in marine sediments on continental margins, and thus preserved over geological timescales. Indeed, our estimated global riverine PBC flux amounts to 20% of the terrestrial organic carbon stored annually in ocean sediments ⁵⁰, suggesting that processes of BC production, protracted storage in terrestrial reservoirs, mobilization and burial in marine sediments thus represent an important geologic atmospheric CO₂ sink.

These findings have implications for our understanding of the role of BC cycling in the face of direct (e.g., land-use) and indirect (climate) anthropogenically-driven change. Some increases in the intensity and frequency of fires with on-going climate change ^{51,52} may enhance BC production. Here, we find river PBC is efficiently exported and stored in sediments rather than degraded to CO₂ en-route to burial in the ocean, suggesting a negative feedback to increased biomass burning. Such interpretations

are likely overly simplistic, by not taking into account seasonal differences between production and erosion of BC. However, the strong correlations imply that these trends are robust at a global scale. Nevertheless, it is clear that further consideration of BC dynamics within river basins at regional and global scales is warranted in order to better constrain this important component of the carbon cycle.

Acknowledgments and author contributions

We thank Michael Hilf for his technical support and laboratory assistance. We also thank Daniel Montlucon for collecting Mississippi River samples, Nicholas Drenzek for assistance in collection of the Eel River samples, Ana Lima, John King and Chris Reddy for collection of Pettaquamscutt River samples, members of the Woods Hole Research Center for collection of Colville and Yukon River samples, Derek Vance and Brandi Revels for providing the logistical foundation for the collection of the Amazon River samples, Dagmar Brandova for her technical assistance, and the ETH Ion Beam Physics Lab AMS Lab colleagues for AMS support. We thank Gregory J. Fiske (Woods Hole Research Center) and Nadja Baltensweiler (University of Zurich Visual Communications Center) for visual support with Figure 3, and 1, respectively. We thank Iselin Medhaug for comments on the manuscript. A. C. acknowledges financial support from the University of Zurich Forschungskredit Fellowship and the Science Faculty of the University of Zurich (No. STWF-18-026). M.Z. acknowledges support from the National Natural Science Foundation of China (No. 41521064). T.E. acknowledges support from the Swiss National Science Foundation (“CAPS-LOCK” and “CAPS-LOCK2” #200021_140850). V.G. acknowledges financial support from an Independent Study Award from the Woods Hole Oceanographic Institution

All authors contributed scientific comments and input on this manuscript. T.E., M.S., D.W. and A.C. contributed to the design of the study. A.C. and D.W. measured the samples for BC content and radiocarbon values. U.H., N.H. and L.W. provided analytical assistance to the radiocarbon measurement of samples and quality control. V.G., A.C., M.S., E.S. and T.E. contributed to data interpretation. A.C. wrote the paper

273 and built the figures. Samples and supplementary data were provided by T.E., G.N,
274 M.U., T.B., C.F., E.S., M.Z., B.V., V.G., M.R. and B.P.E.

275

276

277

278

279

280

281

282

283

284

285

286





Figure 1. Global schematic synthesis of the BC cycle in major reservoirs. Estimates are derived from Supplementary Table 3. The relative size of the reservoir (Tg) is given by the size of the circle. Schematic BC $\Delta^{14}\text{C}$ values are given by the shade of white (modern, post 1950) to dark red (ancient -1000‰) in (circle) reservoirs and river BC (arrow) pools. Fluxes are in Tg yr⁻¹ given by the relative size of the blue and red arrows.



300

301 Figure 2. PBC fluxes (values in Gg yr⁻¹) and PBC $\Delta^{14}\text{C}$ values (in ‰). PBC $\Delta^{14}\text{C}$
 302 values are given by the shade of yellow (bomb) to dark red (ancient -1000‰) for the
 303 river catchment. Rivers were sampled at their outlets. Small colored circles indicate
 304 small mountainous rivers with drainage basin areas less than 250,000 km². The
 305 histogram represents fluxes per river (Gg yr⁻¹), where the grey bar represents the flux
 306 of the Padma, which is the combined fluxes of the Brahmaputra and Ganges rivers.

307

308

309

310

311

312




Figure 3. a) Relationship between PBC yield and suspended sediment yield. PBC was normalized to the biospheric OC yield given in ²⁸. The regression line is $Y_{\text{PBC yield}} = 0.005(Y_{\text{sed}})^{0.69}$; $r^2 = 0.61$; $P < 0.001$. b) Relationship between $\Delta^{14}\text{C}$ values of PBC and POC. The regression line is $Y_{\text{PBC14C}} = 1.05(X_{\text{biosphericOC}}) - 132$; $r^2 = 0.48$; $P < 0.001$.

References

- 1 Foster, G. L., Royer, D. L. & Lunt, D. J. Future climate forcing potentially without precedent in the last 420 million years. *Nature Communications* **8**, 14845, doi:10.1038/ncomms14845
<https://www.nature.com/articles/ncomms14845#supplementary-information> (2017).
- 2 Zimmerman, A. R. & Mitra, S. Trial by Fire: On the Terminology and Methods Used in Pyrogenic Organic Carbon Research. *Frontiers in Earth Science* **5**, doi:10.3389/feart.2017.00095 (2017).
- 3 Santin, C. *et al.* Towards a global assessment of pyrogenic carbon from vegetation fires. *Global Change Biology* **22**, 76-91, doi:10.1111/gcb.12985 (2016).
- 4 Bird, M. I., Wynn, J. G., Saiz, G., Wurster, C. M. & McBeath, A. in *Annual Review of Earth and Planetary Sciences, Vol 43* Vol. 43 *Annual Review of Earth and Planetary Sciences* (eds R. Jeanloz & K. H. Freeman) 273-298 (2015).
- 5 Singh, N., Abiven, S., Torn, M. S. & Schmidt, M. W. I. Fire-derived organic carbon in soil turns over on a centennial scale. *Biogeosciences* **9**, 2847-2857, doi:10.5194/bg-9-2847-2012 (2012).
- 6 Santin, C., Doerr, S. H., Preston, C. M. & Gonzalez-Rodriguez, G. Pyrogenic organic matter production from wildfires: a missing sink in the global carbon cycle. *Global Change Biology* **21**, 1621-1633, doi:10.1111/gcb.12800 (2015).
- 7 Lehmann, J. *et al.* Australian climate-carbon cycle feedback reduced by soil black carbon. *Nature Geoscience* **1**, 832-835, doi:10.1038/ngeo358 (2008).
- 8 Ascough, P. L. *et al.* Variability in oxidative degradation of charcoal: Influence of production conditions and environmental exposure. *Geochimica Et Cosmochimica Acta* **75**, 2361-2378, doi:10.1016/j.gca.2011.02.002 (2011).
- 9 Pingree, M. R. A. & DeLuca, T. H. Function of Wildfire-Deposited Pyrogenic Carbon in Terrestrial Ecosystems. *Frontiers in Environmental Science* **5**, doi:10.3389/fenvs.2017.00053 (2017).
- 10 Ohlson, M., Dahlberg, B., Økland, T., Brown, K. J. & Halvorsen, R. The charcoal carbon pool in boreal forest soils. *Nature Geoscience* **2**, 692, doi:10.1038/ngeo617
<https://www.nature.com/articles/ngeo617#supplementary-information> (2009).
- 11 Aufdenkampe, A. K. *et al.* Riverine coupling of biogeochemical cycles between land, oceans, and atmosphere. *Frontiers in Ecology and the Environment* **9**, 53-60, doi:10.1890/100014 (2011).
- 12 Battin, T. J. *et al.* The boundless carbon cycle. *Nature Geosci* **2**, 598-600 (2009).

- 365 13 Raymond, P. A. *et al.* Global carbon dioxide emissions from inland waters.
366 *Nature* **503**, 355-359, doi:10.1038/nature12760
367 [http://www.nature.com/nature/journal/v503/n7476/abs/nature12760.html#supplement](http://www.nature.com/nature/journal/v503/n7476/abs/nature12760.html#supplementary-information)
368 [ary-information](http://www.nature.com/nature/journal/v503/n7476/abs/nature12760.html#supplementary-information) (2013).
- 369 14 Cole, J. J. *et al.* Plumbing the global carbon cycle: Integrating inland waters
370 into the terrestrial carbon budget. *Ecosystems* **10**, 171-184,
371 doi:10.1007/s10021-006-9013-8 (2007).
- 372 15 Tranvik, L. J. *et al.* Lakes and reservoirs as regulators of carbon cycling and
373 climate. *Limnology and Oceanography* **54**, 2298-2314,
374 doi:10.4319/lo.2009.54.6_part_2.2298 (2009).
- 375 16 Wang, X., Xu, C., Druffel, E. M., Xue, Y. & Qi, Y. Two black carbon pools
376 transported by the Changjiang and Huanghe Rivers in China. *Global*
377 *Biogeochemical Cycles*, n/a-n/a, doi:10.1002/2016GB005509 (2016).
- 378 17 Dittmar, T. *et al.* Continuous flux of dissolved black carbon from a vanished
379 tropical forest biome. *Nature Geoscience* **5**, 618-622, doi:10.1038/ngeo1541
380 (2012).
- 381 18 Jaffe, R. *et al.* Global Charcoal Mobilization from Soils via Dissolution and
382 Riverine Transport to the Oceans. *Science* **340**, 345-347,
383 doi:10.1126/science.1231476 (2013).
- 384 19 Coppola, A. I. & Druffel, E. R. M. Cycling of black carbon in the ocean.
385 *Geophysical Research Letters* **43**, 4477-4482, doi:10.1002/2016GL068574
386 (2016).
- 387 20 Ziolkowski, L. A. & Druffel, E. R. M. Aged black carbon identified in marine
388 dissolved organic carbon. *Geophysical Research Letters* **37**,
389 doi:10.1029/2010gl043963 (2010).
- 390 21 Masiello, C. A. New directions in black carbon organic geochemistry. *Marine*
391 *Chemistry* **92**, 201-213, doi:10.1016/j.marchem.2004.06.043 (2004).
- 392 22 Bird, M. I., Wynn, J. G., Saiz, G., Wurster, C. M. & McBeath, A. The Pyrogenic
393 Carbon Cycle. *Annual Review of Earth and Planetary Sciences* **43**,
394 150223150959000, doi:10.1146/annurev-earth-060614-105038 (2014).
- 395 23 Santín, C. *et al.* Towards a global assessment of pyrogenic carbon from
396 vegetation fires. *Glob Chang Biol*, doi:10.1111/gcb.12985 (2015).
- 397 24 Guerená, D. T. *et al.* Terrestrial pyrogenic carbon export to fluvial
398 ecosystems: Lessons learned from the White Nile watershed of East Africa.
399 *Global Biogeochemical Cycles* **29**, 1911-1928, doi:10.1002/2015gb005095
400 (2015).
- 401 25 Regnier, P. *et al.* Anthropogenic perturbation of the carbon fluxes from land to
402 ocean. *Nature Geosci* **6**, 597-607, doi:10.1038/ngeo1830
403 [http://www.nature.com/ngeo/journal/v6/n8/abs/ngeo1830.html#supplementary-](http://www.nature.com/ngeo/journal/v6/n8/abs/ngeo1830.html#supplementary-information)
404 [information](http://www.nature.com/ngeo/journal/v6/n8/abs/ngeo1830.html#supplementary-information) (2013).

- 405 26 Vorosmarty, C. J. *et al.* Global threats to human water security and river
406 biodiversity. *Nature* **467**, 555-561,
407 doi:[http://www.nature.com/nature/journal/v467/n7315/abs/nature09440.html#s](http://www.nature.com/nature/journal/v467/n7315/abs/nature09440.html#supplementary-information)
408 [upplementary-information](http://www.nature.com/nature/journal/v467/n7315/abs/nature09440.html#supplementary-information) (2010).
- 409 27 Wheatcroft, R. A., Goni, M. A., Hatten, J. A., Pasternack, G. B. & Warrick, J.
410 A. The role of effective discharge in the ocean delivery of particulate organic
411 carbon by small, mountainous river systems. *Limnology and Oceanography*
412 **55**, 161-171, doi:10.4319/lo.2010.55.1.0161 (2010).
- 413 28 Galy, V., Peucker-Ehrenbrink, B. & Eglinton, T. Global carbon export from the
414 terrestrial biosphere controlled by erosion. *Nature* **521**, 204-207,
415 doi:10.1038/nature14400
416 [http://www.nature.com/nature/journal/v521/n7551/abs/nature14400.html#supplement](http://www.nature.com/nature/journal/v521/n7551/abs/nature14400.html#supplementary-information)
417 [ary-information](http://www.nature.com/nature/journal/v521/n7551/abs/nature14400.html#supplementary-information) (2015).
- 418 29 Wiedemeier, D. B. *et al.* Characterization, Quantification and Compound-
419 specific Isotopic Analysis of Pyrogenic Carbon Using Benzene Polycarboxylic
420 Acids (BPCA). *Jove-Journal of Visualized Experiments*, doi:10.3791/53922
421 (2016).
- 422 30 Myers-Pigg, A. N. *et al.* Labile pyrogenic dissolved organic carbon in major
423 Siberian Arctic rivers: Implications for wildfire-stream metabolic linkages.
424 *Geophysical Research Letters* **42**, 377-385, doi:10.1002/2014gl062762
425 (2015).
- 426 31 Lang, S. Q., Fröh-Green, G. L., Bernasconi, S. M. & Wacker, L. Isotopic
427 ($\delta^{13}\text{C}$, $\Delta^{14}\text{C}$) analysis of organic acids in marine samples using wet chemical
428 oxidation. *Limnology and Oceanography: Methods* **11**, 161-175,
429 doi:10.4319/lom.2013.11.161 (2013).
- 430 32 Reisser, M., Purves, R. S., Schmidt, M. W. I. & Abiven, S. Pyrogenic Carbon
431 in Soils: A Literature-Based Inventory and a Global Estimation of Its Content
432 in Soil Organic Carbon and Stocks. *Frontiers in Earth Science* **4**,
433 doi:10.3389/feart.2016.00080 (2016).
- 434 33 Galy, V. & Eglinton, T. Protracted storage of biospheric carbon in the Ganges-
435 Brahmaputra basin. *Nature Geosci* **4**, 843-847,
436 doi:[http://www.nature.com/ngeo/journal/v4/n12/abs/ngeo1293.html#suppleme](http://www.nature.com/ngeo/journal/v4/n12/abs/ngeo1293.html#supplementary-information)
437 [ntary-information](http://www.nature.com/ngeo/journal/v4/n12/abs/ngeo1293.html#supplementary-information) (2011).
- 438 34 Burdige, D. J. Preservation of Organic Matter in Marine Sediments: Controls,
439 Mechanisms, and an Imbalance in Sediment Organic Carbon Budgets?
440 *Chemical Reviews* **107**, 467-485, doi:10.1021/cr050347q (2007).
- 441 35 Hanke, U. M. *et al.* What on Earth have we been burning? Deciphering
442 sedimentary records of pyrogenic carbon. *Environmental Science &*
443 *Technology*, doi:10.1021/acs.est.7b03243 (2017).

- 444 36 Marwick, T. R. *et al.* The age of river-transported carbon: A global
445 perspective. *Global Biogeochemical Cycles* **29**, 122-137,
446 doi:10.1002/2014GB004911 (2015).
- 447 37 Cotrufo, M. F. *et al.* Redistribution of pyrogenic carbon from hillslopes to
448 stream corridors following a large montane wildfire. *Global Biogeochemical*
449 *Cycles* **30**, 1348-1355, doi:10.1002/2016GB005467 (2016).
- 450 38 Frueh, W. T. & Lancaster, S. T. Correction of deposit ages for inherited ages
451 of charcoal: implications for sediment dynamics inferred from random
452 sampling of deposits on headwater valley floors. *Quaternary Science Reviews*
453 **88**, 110-124, doi:<http://dx.doi.org/10.1016/j.quascirev.2013.10.029> (2014).
- 454 39 Tao, S., Eglinton, T. I., Montluçon, D. B., McIntyre, C. & Zhao, M. Pre-aged
455 soil organic carbon as a major component of the Yellow River suspended
456 load: Regional significance and global relevance. *Earth and Planetary*
457 *Science Letters* **414**, 77-86, doi:<https://doi.org/10.1016/j.epsl.2015.01.004>
458 (2015).
- 459 40 Feng, X. *et al.* Differential mobilization of terrestrial carbon pools in Eurasian
460 Arctic river basins. *Proceedings of the National Academy of Sciences* **110**,
461 14168-14173, doi:10.1073/pnas.1307031110 (2013).
- 462 41 Schefuß, E. *et al.* Hydrologic control of carbon cycling and aged carbon
463 discharge in the Congo River basin. *Nature Geoscience* **9**, 687,
464 doi:10.1038/ngeo2778
465 <https://www.nature.com/articles/ngeo2778#supplementary-information> (2016).
- 466 42 Santin, C. *et al.* Towards a global assessment of pyrogenic carbon from
467 vegetation fires. *Global Change Biology* **22**, 76-91, doi:10.1111/gcb.12985
468 (2016).
- 469 43 Regnier, P. *et al.* Anthropogenic perturbation of the carbon fluxes from land to
470 ocean. *Nature Geoscience* **6**, 597-607, doi:10.1038/ngeo1830 (2013).
- 471 44 Marin-Spiotta, E. *et al.* Long-term stabilization of deep soil carbon by fire and
472 burial during early Holocene climate change. *Nature Geosci* **7**, 428-432,
473 doi:10.1038/ngeo2169
474 [http://www.nature.com/ngeo/journal/v7/n6/abs/ngeo2169.html#supplementary-](http://www.nature.com/ngeo/journal/v7/n6/abs/ngeo2169.html#supplementary-information)
475 [information](http://www.nature.com/ngeo/journal/v7/n6/abs/ngeo2169.html#supplementary-information) (2014).
- 476 45 Wagner, S., Cawley, K. M., Rosario-Ortiz, F. L. & Jaffe, R. In-stream sources
477 and links between particulate and dissolved black carbon following a wildfire.
478 *Biogeochemistry* **124**, 145-161, doi:10.1007/s10533-015-0088-1 (2015).
- 479 46 Schmidt, M. W. I. *et al.* Persistence of soil organic matter as an ecosystem
480 property. *Nature* **478**, 49-56 (2011).
- 481 47 Lehmann, J. *et al.* *Persistence of biochar in soil.* (2015).
- 482 48 Amelung, W., Brodowski, S., Sandhage-Hofmann, A. & Bol, R. in *Advances in*
483 *Agronomy* Vol. 100 155-250 (Academic Press, 2008).

- 484 49 Hedges, J. I. & Keil, R. G. Sedimentary organic matter preservation: an
485 assessment and speculative synthesis. *Marine Chemistry* **49**, 81-115,
486 doi:[https://doi.org/10.1016/0304-4203\(95\)00008-F](https://doi.org/10.1016/0304-4203(95)00008-F) (1995).
- 487 50 Masiello, C. A. & Druffel, E. R. M. Black carbon in deep-sea sediments.
488 *Science* **280**, 1911-1913, doi:10.1126/science.280.5371.1911 (1998).
- 489 51 Ciais, P., C. Sabine, G. Bala, L. Bopp, V. Brovkin, J. Canadell, A. Chhabra, R.
490 DeFries, J. Galloway, M. Heimann, C. Jones, C. Le Quéré, R.B. Myneni, S.
491 Piao and P. Thornton, . in *Climate Change 2013: The Physical Science Basis.*
492 *Contribution of Working Group I to the Fifth Assessment Report*
493 *of the Intergovernmental Panel on Climate Change* (ed T.F. Stocker, D. Qin,
494 G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex.,
495 P.M. Midgley) (Cambridge University Press, 2013).
- 496 52 Flannigan, M. et al. Global wildland fire season severity in the 21st century.
497 *Forest Ecology and Management* **294**, 54-61,
498 doi:<https://doi.org/10.1016/j.foreco.2012.10.022> (2013).
499